



# Blood Biomarkers of Late Pregnancy Exposure to Trihalomethanes in Drinking Water and Fetal Growth Measures and Gestational Age in a Chinese Cohort

Wen-Cheng Cao, Qiang Zeng, Yan Luo, Hai-Xia Chen,  
Dong-Yue Miao, Li Li, Ying-Hui Cheng, Min Li, Fan Wang,  
Ling You, Yi-Xin Wang, Pan Yang, and Wen-Qing Lu

<http://dx.doi.org/10.1289/ehp.1409234>

Received: 18 September 2014

Accepted: 13 August 2015

Advance Publication: 4 September 2015

This article will be available in a 508-conformant form upon final publication. If you require a 508-conformant version before then, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.



National Institute of  
Environmental Health Sciences

# **Blood Biomarkers of Late Pregnancy Exposure to Trihalomethanes in Drinking Water and Fetal Growth Measures and Gestational Age in a Chinese Cohort**

Wen-Cheng Cao,<sup>1,2</sup> Qiang Zeng,<sup>1,2</sup> Yan Luo,<sup>1,2</sup> Hai-Xia Chen,<sup>1,2</sup> Dong-Yue Miao,<sup>1,2</sup> Li Li,<sup>3</sup>  
Ying-Hui Cheng,<sup>4</sup> Min Li,<sup>1,2</sup> Fan Wang,<sup>1,2</sup> Ling You,<sup>1,2</sup> Yi-Xin Wang,<sup>1,2</sup> Pan Yang,<sup>1,2</sup> and  
Wen-Qing Lu<sup>1,2</sup>

<sup>1</sup>Department of Occupational and Environmental Health, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, PR China; <sup>2</sup>Key Laboratory of Environment and Health, Ministry of Education & Ministry of Environmental Protection, and State Key Laboratory of Environment and Health (incubating), School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, PR China; <sup>3</sup>Department of Gynecology and Obstetrics, Wuhan NO.1 Hospital, Wuhan, Hubei, PR China; <sup>4</sup>Department of Gynecology and Obstetrics, Xiaonan Maternal and Child Care Service Centre, Xiaogan, Hubei, PR China

**Address correspondence to** Wen-Qing Lu, Department of Occupational and Environmental Health, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, PR China. Telephone: +86-27-83610149. E-mail:

[luwq@mails.tjmu.edu.cn](mailto:luwq@mails.tjmu.edu.cn) (W. Q. Lu)

**Running title:** Maternal blood trihalomethanes and fetal growth

**Acknowledgments:** We sincerely thank all staff members in the two hospitals who contributed to this study and all participants for their great cooperation. This research was supported by the Open Funds of State Key Laboratory of Freshwater Ecology and Biotechnology (Grant no: 2012FB08) and the Fundamental Research Funds for the Central Universities of China (HUST: 2013YGYL001).

**Competing financial interests:** The authors declare they have no actual or potential competing financial interests.

## Abstract

**Background:** Previous studies have suggested that elevated exposure to disinfection by-products (DBPs) in drinking water during gestation may result in adverse birth outcomes. However, the findings of these studies remain inconclusive.

**Objective:** The purpose of our study was to examine the association between blood biomarkers of late pregnancy exposure to trihalomethanes (THMs) in drinking water and fetal growth and gestational age.

**Methods:** We recruited 1184 pregnant women between 2011 and 2013 in Wuhan and Xiaogan City, Hubei, China. Maternal blood THM concentrations, including chloroform (TCM), bromodichloromethane (BDCM), dibromochloromethane (DBCM) and bromoform (TBM), were measured as exposure biomarkers during late pregnancy. We estimated associations with gestational age and fetal growth indicators [birth weight, birth length, and small for gestational age (SGA)].

**Results:** Total THMs (TTHMs, sum of TCM, BDCM, DBCM and TBM) were associated with lower mean birth weight (-60.9 g; 95% CI: -116.2, -5.6 for the highest versus lowest tertile;  $p$  for trend = 0.03), and BDCM and DBCM exposures were associated with smaller birth length (e.g., -0.20 cm; 95% CI: -0.37, -0.04 for the highest versus lowest tertile of DBCM;  $p$  for trend = 0.02). SGA was increased in association with the second and third tertiles of TTHMs (OR = 2.91; 95%CI: 1.32, 6.42 and OR = 2.25; 95%CI: 1.01, 5.03;  $p$  for trend = 0.08).

**Conclusions:** Our results suggested that elevated maternal THM exposure may adversely affect fetal growth.

## Introduction

Chlorine, due to its efficacy and cost-effectiveness, has been extensively used in the treatment of drinking water to reduce the risk of waterborne disease worldwide, including in China. However, chlorine and other disinfectants can react with natural organic and inorganic matter that occurs in water to form disinfection by-products (DBPs), which have been suggested to be potentially carcinogenic and to exert reproductive and developmental toxicities (Nieuwenhuijsen et al. 2010). Trihalomethanes (THMs) are the most abundant DBP class in drinking water and include chloroform (TCM), bromodichloromethane (BDCM), dibromochloromethane (DBCM) and bromoform (TBM) (Nieuwenhuijsen et al. 2000). Widespread exposure to THMs can result from ingestion, inhalation and dermal absorption during routine water-use activities such as drinking, washing, showering, bathing and swimming. Based on the potential adverse health effects of exposure to DBPs, four THMs have been regulated in the European Union, USA and other countries (e.g., in Australia, and China).

Toxicological studies have found that THMs may result in adverse reproductive effects. Exposure to TCM, DBCM through oral administration has been shown to cause fetal toxicity in rats including decreased body weight, body length and survival rate (Ruddick et al. 1983). Exposure to BDCM at high doses has also been shown to cause pregnancy loss in rats (Bielmeier et al. 2001). A number of epidemiological studies have also examined the relationship between DBP exposure and adverse reproductive outcomes, including pregnancy loss, birth defects and fetal growth restriction (Dodds et al. 2004; Grazuleviciene et al. 2013; Hoffman et al. 2008b; Infante-Rivard 2004; Savitz et al. 2006; Toledano et al. 2005); however, the findings of these studies are equivocal. One of the major limitations of previous studies is inaccurate exposure assessment (Nieuwenhuijsen et al. 2009).

Most researchers took advantage of routinely collected measurements of THM concentrations in public water supplies as a surrogate of exposure, and some also combined data from water-use activities to estimate internal THM dose. However, these exposure assessments may result in misclassification of exposure by several factors: spatial and temporal variability of THMs in water systems, the contribution of different exposure routes, inter- and intra-individual variability in water usage (including residential mobility), and inter-and intra-individual physiological differences in absorption, distribution, metabolism and excretion of the four THMs (Backer et al. 2000, 2008; Leavens et al. 2007).

Exposure biomarkers can represent integrative measures of all routes of exposure and provide an accurate exposure assessment for specific exposure windows. THM concentrations in blood and alveolar air samples have been measured to assess internal dose levels of THMs (Gordon et al. 2006; Lakind et al. 2009). Although the collection of breath samples is non-invasive, THM concentrations were generally undetectable before high levels of exposure (Weisel et al. 1999). In contrast, blood THM concentrations were generally more sensitive to low levels of exposure (Backer et al. 2000; Weisel et al. 1999). Although the elimination half-life of THMs in blood is short (minutes-hours), there are believed to be steady-state concentrations due to repeated and relatively consistent exposure to THMs (Blount et al. 2011). Several factors have been associated with blood THM levels including THM concentrations in water distribution systems, water-use activities (e.g., bathing/showering, and swimming), personal socio-demographic characteristics (e.g., age, body mass index, education, and household income) and genetic and physiological differences (Aggazzotti et al. 1998; Backer et al. 2008; Caro and Gallego 2007; Lynberg et al. 2001; Nuckols et al. 2005; Riederer et al. 2014; Rivera-Nunez et al. 2012; Zeng et al. 2014b).

We conducted a study in Wuhan and Xiaogan city, Hubei, China to investigate the relationships between exposure to drinking water DBPs and birth outcomes. In our study, whole blood THM levels were determined to assess the internal dose of THM exposure. To our knowledge, our study is the first to use THM levels in whole blood as exposure biomarkers to evaluate the effects of exposure to THMs in drinking water on birth outcomes.

## **Methods**

### **Study participants**

We conducted a study in two contiguous cities, Wuhan and Xiaogan, Hubei, China. The water distribution systems in the two cities are supplied by surface water sources, and chlorine is used in water treatment process. Women in late pregnancy during July 2011 to July 2012 in Wuhan and during October 2012 to December 2013 in Xiaogan were invited to participate in the study. The study was approved by the Ethics Committee of Tongji Medical College, and all participants provided written informed consent at the time of enrollment.

During the study period, a total of 997 and 750 pregnant women during late pregnancy ( $\geq 35$  weeks) in Wuhan and Xiaogan agreed to participate in the study, respectively. Of them, 1261 (72%) provided blood samples for analysis. We restricted our analyses to single gestation live infants, whose mothers lived in the local city for at least 1 year ( $n = 77$  excluded participants), resulting in a total of 1184 births.

### **Questionnaires**

All participants took part in a face-to-face interview conducted by the trained investigators to complete a structured questionnaire on the first day of hospital admittance waiting for delivery. The

questionnaire included demographics, lifestyle, occupational exposures during pregnancy, gravidity, parity, case history, and routine water-use activities. Data regarding water-use activities included source of drinking water, use of boiled-water and filtered water, the total volume of tap-water consumption per day (number multiplied by glass size), minutes of showering/bathing per week (frequency  $\times$  duration of bathing/showering), minutes per week spent washing dishes and clothes by hand without gloves, respectively (frequency  $\times$  duration of each activity) and swimming pool attendance (yes/no) during pregnancy.

### **Outcome data**

Basic information regarding the infants, including gestational age, gender, birth length and birth weight, were collected from the clinical birth records. Gestational age was based on the interval between the last menstrual period and the date of delivery of the infant. Small for gestational age (SGA) was defined as a live-born infant below the 10<sup>th</sup> percentile of birth weight for gestational age in a national Chinese referent population (Chen and Jin 2011).

### **Blood sample collection and blood THM analyses**

A 5-mL blood sample was collected by nurses  $\geq$  2 hours after last showering/bathing on the first day of hospital admittance of pregnant women waiting for delivery. After the blood draw, the tubes were shaken to dissolve the anticoagulant (potassium-ethylenediaminetetraacetic acid) immediately, kept in coolers and then shipped to the laboratory. The blood samples were kept at 4 °C before they were analyzed for THMs within 2 weeks (Bonin et al. 2005).

Concentrations of THMs in blood samples were determined by headspace solid phase micro-extraction (SPME)-gas chromatography with an electron capture detector (GC/ECD, Agilent



Technologies 6890 N). The detailed method and quality control have been described in our previous study (Zeng et al. 2013). Briefly, we sealed 3-mL blood samples in 10-mL headspace vials. We then heated (20 °C) and agitated (300 rpm) samples using a magnetic stirrer to facilitate extraction of volatiles from the sample headspace onto an SPME fiber. After extraction, we immediately inserted the fiber into a hot GC inlet and maintained it for 3 minutes. We identified the four individual THMs according to retention times. Final quantification was based on procedural standard calibration curves. The limit of detection (LOD) for TCM, BDCM, DBCM and TBM were 1.9, 0.5, 0.7 and 2.0ng/L, respectively. Concentrations below the LOD were assigned with LOD/ $\sqrt{2}$  for the analysis.

### **Statistical methods**

The Predictive Analytics Suite Workstation (PASW) version 18.0 (IBM corporation, Armonk, New York, USA) was used for the analysis. Descriptive statistics for demographics, birth outcomes, and maternal blood THMs were conducted. To compare differences of fetal growth measures and gestational age in all categorical variables, parametric and nonparametric methods were appropriately used to test statistical significance. In addition, Pearson correlation analysis was used to examine the association between maternal age and fetal growth measures. Br-THM concentration was defined as the sum of BDCM, DBCM and TBM in blood. Total THM (TTHM) concentration was defined as the sum of TCM and Br-THMs in blood. Because the detectable percentage of blood BDCM, DBCM and TBM concentrations is not high, we used a three-level variable to categorize participants into low exposure (<LOD) and equally-sized medium and high exposure groups. We divided maternal blood TCM, Br-THM, and TTHM concentrations into tertiles based on measured

values (none was <LOD), and used the lowest level as the reference. We conducted tests for trend by treating the blood THMs as an ordinal categorical variable in regression models.

General linear models were applied to analyze the association between maternal blood THM level and fetal growth and gestational duration indices (birth weight, birth length and gestational age).

Logistic regression models were used to estimate odds ratios (ORs) and 95 percent confidence intervals (95% CIs) for SGA infants. Covariates were included in the final models for each fetal outcome based on biological and statistical considerations. For the statistical consideration, potential confounders [gestational age, prenatal body mass index (BMI), weight gain during pregnancy, infant's gender, study city, education, and household income] were entered into the final multivariable model with  $p$ -value <0.2 for unadjusted associations with fetal outcomes. Maternal age and parity were included in final models based on biological consideration, as previous studies have suggested that they are predictors of fetal growth (Da Silva. 2012; Shah. 2010).

All regression models were adjusted for the following dichotomous variables: prenatal BMI (<28/≥28 kg/m<sup>2</sup>), weight gain during pregnancy (<15/≥15 kg), infant's gender (male/female), parity (no child/≥one child), and study city (Xiaogan/Wuhan). As a continuous variable, maternal age (squared, years<sup>2</sup>) was included in all regression models; gestational age (weeks) was only entered into the models for birth weight and birth length. As a categorical variable, education (less than primary school, junior and senior high school, college and above) was included in the models for birth weight, birth length and SGA; household income (<3000, 3000-<5000, ≥5000 Yuan) was included in the models for gestational age and SGA. Statistical significance was defined as a  $p$  value <0.05. And statistically suggestive was defined as a  $p$  value <0.10 (Zeng et al. 2013).

## Results

### Characteristics of the study population

The demographic characteristics of mothers and their infants are summarized in Table 1. Of the 1184 single gestation live births, 60 (5.1%) were classified as SGA. The mean birth weight, birth length and gestational age were  $3322 \pm 404$  g,  $50 \pm 1$  cm and  $39 \pm 5$  weeks, respectively. Mean maternal age was  $29 \pm 5$  years old. The majority of mothers had prenatal BMI  $<28$  kg/m<sup>2</sup> (68.5%), gained  $>15$  g during pregnancy (61.6%), reported drinking  $<1200$  ml of water per day (55.8%), and did not swim during pregnancy (99.4%), or hand wash dishes (73.0%) or clothes (53.4%) without gloves during pregnancy.

Water-use activities were not significantly associated with birth outcomes, with the exception of tap water consumption and gestational age (mean values of  $39.2 \pm 1.3$  and  $39.0 \pm 1.3$  weeks among mothers who reported drinking  $<1200$  and  $\geq 1200$  mL/day, respectively,  $p$ -value  $<0.01$ ) (Table 1). However, the association was only suggestive after adjusting confounding ( $-0.15$  weeks shorter; 95% CI:  $-0.31, 0.01$  for  $\geq 1200$  versus  $<1200$  mL/day;  $p$ -value = 0.07).

### Maternal blood THM concentrations

The distribution of maternal blood THM concentrations among the study participants is presented in Table 2. TCM was detected in 92.5% of the blood samples, whereas BDCM, DBCM and TBM were found at a lower frequency, ranging from 22.6% to 57.4%. The geometric mean (median) concentrations of TCM, Br-THMs and TTHMs were 40.7 (50.7) ng/L, 5.3 (5.6) ng/L and 52.3 (57.7) ng/L, respectively.

## Maternal blood THMs and fetal growth

Table 3 presents regression coefficients [ $\beta$  (95% CI)] for fetal growth associated with categories of maternal blood THM concentrations. We found no statistically significant associations between maternal blood THM concentrations and gestational age. Maternal TTHM concentrations in the second and third tertiles (44.2 - 74.4 and >74.4 ng/L, respectively) were associated with lower birth weight relative to the lowest tertile (<44.2 ng/L), with estimated mean differences of -59.09 g (95%CI: -114.46, -3.71) and -60.88 g (95%CI: -116.18, -5.58), respectively ( $p$  for trend = 0.03). Additionally, there was a suggestive negative association between birth weight and TCM (-48.23 g; 95% CI:-103.64, 7.19 for the third versus first tertile,  $p$  for trend = 0.08). BDCM and DBCM were negatively associated with length at birth, with estimated mean decreases of 0.15 cm (95%CI: -0.29, -0.01) and 0.20 cm (95%CI: -0.37, -0.04), respectively, for the highest versus lowest exposure groups ( $p$  for trend of 0.04 and 0.02, respectively).

The ORs and 95%CI for SGA and maternal blood THM concentrations are shown in Table 4. Exposure to Br-THMs was positively but only suggestive associated with SGA (OR = 1.48; 95% CI: 0.71, 3.04 and OR = 1.92; 95%CI: 0.98, 3.79 for the 2<sup>nd</sup> and 3<sup>rd</sup> exposure groups, respectively;  $p$  for trend = 0.06). SGA was significantly increased in association with the 2<sup>nd</sup> and 3<sup>rd</sup> tertiles of TTHMs (OR = 2.91; 95%CI: 1.32, 6.42 and OR = 2.25; 95%CI: 1.01, 5.03, respectively,  $p$  for trend = 0.08).

## Discussion

We determined the maternal blood THM concentrations as an internal dose level of THM exposure, which could represent an accurate and integrative measure of all routes and sources of exposure. Because blood concentrations are strongly influenced by very recent exposure, and showering and

bathing have been shown to have a stronger influence on blood levels than other water-use activities (Nuckols et al. 2005), we collected blood samples after at least 2 hours since last showering/bathing to gain a relatively steady state of THMs in the blood. Two studies have shown that the blood samples were taken after 30 minutes since last showering/bathing can provide a window to a steady-state level (Ashley et al. 2005; Silva et al. 2013). Ashley et al (2005) reported decrease in blood THM concentrations from 5 min to 30 min post shower/bath among 7 young and healthy subjects. Silva et al (2013) also found that blood THM concentrations dropped rapidly during the first 30 minutes after showering among 100 study participants following a controlled showering exposure. Consistent with two previous reports, TCM was the main component of blood TTHMs in our study population (>70%) (Miles et al. 2002; Riederer et al. 2014). The median concentrations of blood THM were higher than reported for a representative sample of US adults (NHANES participants in 1999–2006) (Riederer et al. 2014) and a group of 150 US women (Rivera-Nunez et al. 2012), but were similar to levels reported for 401 men from Wuhan, China (Zeng et al. 2013).

TTHMs were associated with a significant decrease in mean birth weight, with similar estimated reductions for the 2<sup>nd</sup> and 3<sup>rd</sup> tertiles of exposure compared with the 1<sup>st</sup>. We also found some evidence for associations between exposure to TTHMs during late pregnancy and the risk of SGA, which was consistent with previous studies that reported a small increased risk of SGA for high exposure TTHMs (Hinckley et al. 2005; Hoffman et al. 2008b; Porter et al. 2005; Wright et al. 2003, 2004). Many previous studies have characterized exposures based on total blood THM concentrations, though a few have evaluated exposures to individual THMs. For the individual THMs in our study, we found associations of reduced birth length with individual brominated THMs (e.g., BDCM and DBCM) but not with TCM, however, we found a suggestive association of

reduced birth weight with TCM but not with individual brominated THMs, which was consistent with previous toxicological studies in rats showing that brominated THMs were more harmful to the fetus than TCM (Narotsky et al. 1997; Ruddick et al. 1983). Exposure to brominated THMs (e.g., BDCM, 50 and 75 mg/kg; DBCM, 200 mg/kg) can result in reduced body length and increased rate of fetal resorptions, while only exposure to higher dose of TCM (400mg/kg) can result in reduced fetal body weight (Narotsky et al. 1997; Ruddick et al. 1983). With the exception of our study, only one other study (Patelarou et al. 2011) has reported an excess risk of SGA based on length (a live-born infant below the 10<sup>th</sup> percentile of birth length for gestational age in a referent population) for the higher tertile of brominated THMs (adjusted OR = 1.3, 95%CI, 0.5 to 2.7). The association between THMs and birth length requires further investigations.

We did not find a statistically significant association between gestational age and THM exposure during pregnancy, which was also consistent with the results of several other studies, including a recent meta-analysis (Aggazzotti et al. 2004; Grellier et al. 2010; Jaakkola et al. 2001). However, others found that exposure to high TTHMs may prolong gestational duration and reduce the risk of preterm delivery (Hoffman et al. 2008a; Lewis et al. 2007; Wright et al. 2004). These inconsistent results may be attributed to differences in the characteristics of the study population, in exposure assessments, and in the ability of controlling for confounding factors.

Several limitations of our study should be mentioned. First, because the demographic profiles and the neonatal infant characteristics were different between our studied cities, we adjusted for potential risk factors for fetal growth that varied between cities to control for potential confounding between cities. Although some of covariates (e.g., season, smoking, second-hand smoking, alcohol use, gravidity and maternal medical risk factors during pregnancy) varied between cities, they were

not included in the final models because they did not predict fetal outcomes with  $p < 0.2$  in bivariable models. Nevertheless, the study city was a predictor of gestational age. This suggests that there may be other unmeasured factors that affect gestational duration and vary by study city (Hoffman et al. 2008a). In addition, the detailed data on other potential confounders were missing, including prenatal care access, isolated maternal medical risk factors (e.g., chronic nephritic syndrome, anemia, uterine bleeding during pregnancy), and dietary habits (e.g., fasting), which might bias the relationships observed in our study. Furthermore, our previous study conducted in a water supply system in Wuhan has shown that THM levels in drinking water are below the regulatory limits of China (Zeng et al. 2014a). However, THM levels in the other water supply systems in Wuhan, as well as in Xiaogan, were missing. Thus, the contributions of water THMs to the internal dose of THMs and birth outcomes were unclear.

Second, we relied on a single blood sample during the third trimester to estimate exposure. Although it has been suggested that blood concentrations at a single point in time may reflect steady-state levels (Blount et al. 2011), high exposure events such as showering, bathing, and swimming, can have a substantial effect on blood concentrations (Nuckols et al. 2005; Silva et al. 2013). Changes in routine water-use activities in late pregnancy, and dietary changes (e.g., fasting) before delivery, also may cause blood THM concentrations to fluctuate, both within and between days (Ashley et al. 2005; Riederer et al. 2014). Thus, future studies should collect multiple blood samples to provide a more accurate measure of steady-state levels, and use longer-lived exposure biomarkers (e.g., protein or DNA adducts) to avoid this limitation (Blount et al. 2011).

Third, although the third trimester is the most important for fetal growth (Diamond 2001; Grellier et al. 2010), some studies have found an association of TLBW (term low birth weight) or SGA with

high levels of TTHMs during the second trimester (Lewis et al. 2006; Wright et al. 2003), suggesting that exposure prior to the third trimester may also hamper fetal growth. Pharmacokinetic changes during pregnancy, including increases in plasma volume, changes in blood protein binding, and fat accumulation during the first two trimesters; and increased CYP1A1 and CYP2D6 activity in third trimester; also may influence the relationship between environmental exposures and resulting blood THM concentrations (Anderson 2005; Choi et al. 2013). Therefore, assessment of exposure with exposure biomarkers during different trimesters of pregnancy deserves attention in future studies.

Finally, we estimated associations between exposure to drinking water DBPs and fetal growth based on blood THM concentrations. However, people are generally exposed to DBP mixtures in drinking water that may include other DBPs that are more toxic to fetal growth than THMs (Richardson et al. 2007). For example, previous studies have reported that SGA and decreased birth weight are associated with urine TCAA, a biomarker that reflects ingestion of DBPs in chlorinated drinking water (Costet et al. 2012; Zhou et al. 2012). Because the physicochemical properties, exposure route, metabolism and toxicity among different DBP classes vary, THMs may not be a valid marker of exposure to other DBPs that may be more etiologically relevant (Zeng et al. 2014b). Thus, the associations of other specific DBPs should be considered in future studies.

## **Conclusions**

In the present study, we used whole blood THMs as exposure biomarkers to estimate associations between exposure to THMs in drinking water and fetal growth outcomes and gestational age. We found that elevated maternal blood THM concentrations were associated with decreased birth weight, reduced birth length and increased risk of SGA, suggesting that elevated maternal THM



exposure during late pregnancy may adversely affect fetal growth. However, one-time blood THM concentrations may not be good biomarkers of DBP exposure in general during pregnancy. Further studies with different exposure biomarkers for trimester-specific exposure monitoring are needed.

## References

- Aggazzotti G, Fantuzzi G, Righi E, Predieri G. 1998. Blood and breath analyses as biological indicators of exposure to trihalomethanes in indoor swimming pools. *Sci Total Environ* 217(1-2):155-163.
- Aggazzotti G, Righi E, Fantuzzi G, Biasotti B, Ravera G, Kanitz S, et al. 2004. Chlorination by-products (CBPs) in drinking water and adverse pregnancy outcomes in Italy. *J Water Health* 2(4):233-247.
- Anderson DG. 2005. Pregnancy-induced changes in pharmacokinetics: a mechanistic-based approach. *Clin Pharmacokinet* 44:989-1008.
- Ashley DL, Blount BC, Singer PC, Depaz E, Wilkes C, Gordon S, et al. 2005. Changes in blood trihalomethane concentrations resulting from differences in water quality and water use activities. *Arch Environ Occup Health* 60(1):7-15.
- Backer LC, Ashley DL, Bonin MA, Cardinali FL, Kieszak SM, Wooten JV. 2000. Household exposures to drinking water disinfection by-products: whole blood trihalomethane levels. *J Expo Anal Environ Epidemiol* 10(4):321-326.
- Backer LC, Lan Q, Blount BC, Nuckols JR, Branch R, Lyu CW, et al. 2008. Exogenous and endogenous determinants of blood trihalomethane levels after showering. *Environ Health Perspect* 116(1):57-63.
- Bielmeier SR, Best DS, Guidici DL, Narotsky MG. 2001. Pregnancy loss in the rat caused by bromodichloromethane. *Toxicol Sci* 59(2):309-315.
- Blount BC, Backer LC, Aylward LL, Hays SM, LaKind JS. 2011. Human Exposure Assessment for DBPs: Factors Influencing Blood Trihalomethane Levels. In: *Encyclopedia of Environmental Health* (Nriagu JO, Ed). Burlington: Elsevier, 100-107.
- Bonin MA, Silva LK, Smith MM, Ashley DL, Blount BC. 2005. Measurement of trihalomethanes and methyl tert-butyl ether in whole blood using gas chromatography with high-resolution mass spectrometry. *J Anal Toxicol* 29(2):81-89.
- Caro J, Gallego M. 2007. Assessment of exposure of workers and swimmers to trihalomethanes in an indoor swimming pool. *Environ Sci Technol* 41(13):4793-4798.
- Chen C, Jin H. 2011. Neonatal Classification. In: *Practice of Neonatology* (Shao XM, Ye HM, Qiu XS, Eds). Beijing: People's Medical Publishing House, 46-47.
- Choi SY, Koh KH, Jeong H. 2013. Isoform-specific regulation of cytochromes P450 expression by estradiol and progesterone. *Drug Metab Dispos* 41(2):263-269.
- Costet N, Garlantezec R, Monfort C, Rouget F, Gagniere B, Chevrier C, et al. 2012. Environmental and Urinary Markers of Prenatal Exposure to Drinking Water Disinfection By-Products, Fetal

- Growth, and Duration of Gestation in the PELAGIE Birth Cohort (Brittany, France, 2002-2006). *Am J Epidemiol* 175(4):263-275.
- Da Silva TR. 2012. Nonbiological maternal risk factor for low birth weight on Latin America: a systematic review of literature with meta-analysis. *Einstein (Sao Paulo)* 10(3): 380-385.
- Diamond FJ. 2001. Fetal growth programs future health: causes and consequences of intrauterine growth retardation. *Adv Pediatr* 48:245-272.
- Dodds L, King W, Allen AC, Armson BA, Fell DB, Nimrod C. 2004. Trihalomethanes in public water supplies and risk of stillbirth. *Epidemiology* 15(2):179-186.
- Gordon SM, Brinkman MC, Ashley DL, Blount BC, Lyu C, Masters J, et al. 2006. Changes in breath trihalomethane levels resulting from household water-use activities. *Environ Health Perspect* 114(4):514-521.
- Grazuleviciene R, Kapustinskiene V, Vencloviene J, Buinauskiene J, Nieuwenhuijsen MJ. 2013. Risk of congenital anomalies in relation to the uptake of trihalomethane from drinking water during pregnancy, 274-282.
- Grellier J, Bennett J, Patelarou E, Smith RB, Toledano MB, Rushton L, et al. 2010. Exposure to Disinfection By-products, Fetal Growth, and Prematurity. *Epidemiology* 21(3):300-313.
- Hinckley AF, Bachand AM, Reif JS. 2005. Late pregnancy exposures to disinfection by-products and growth-related birth outcomes. *Environ Health Perspect* 113(12): 1808-1813.
- Hoffman CS, Mendola P, Savitz DA, Herring AH, Loomis D, Hartmann KE, et al. 2008a. Drinking water disinfection by-product exposure and duration of gestation. *Epidemiology* 19(5):738-746.
- Hoffman CS, Mendola P, Savitz DA, Herring AH, Loomis D, Hartmann KE, et al. 2008b. Drinking water disinfection by-product exposure and fetal growth. *Epidemiology* 19(5):729-737.
- Infante-Rivard C. 2004. Drinking water contaminants, gene polymorphisms, and fetal growth. *Environ Health Perspect* 112(11):1213-1216.
- Jaakkola JJ, Magnus P, Skrandal A, Hwang BF, Becher G, Dybing E. 2001. Foetal growth and duration of gestation relative to water chlorination. *Occup Environ Med* 58(7):437-442.
- Lakind JS, Naiman DQ, Hays SM, Aylward LL, Blount BC. 2009. Public health interpretation of trihalomethane blood levels in the United States: NHANES 1999–2004. *Journal of Exposure Science and Environmental Epidemiology* 20(3):255-262.
- Leavens TL, Blount BC, DeMarini DM, Madden MC, Valentine JL, Case MW, et al. 2007. Disposition of bromodichloromethane in humans following oral and dermal exposure. *Toxicol Sci* 99(2):432-445.

- Lewis C, Suffet IH, Hoggatt K, Ritz B. 2007. Estimated effects of disinfection by-products on preterm birth in a population served by a single water utility. *Environ Health Perspect* 115(2):290-295.
- Lewis C, Suffet IH, Ritz B. 2006. Estimated effects of disinfection by-products on birth weight in a population served by a single water utility. *Am J Epidemiol* 163(1):38-47.
- Lynberg M, Nuckols JR, Langlois P, Ashley D, Singer P, Mendola P, et al. 2001. Assessing exposure to disinfection by-products in women of reproductive age living in Corpus Christi, Texas, and Cobb County, Georgia: Descriptive results and methods. *Environ Health Perspect* 109(6):597-604.
- Miles AM, Singer PC, Ashley DL, Lynberg MC, Mendola P, Langlois PH, et al. 2002. Comparison of trihalomethanes in tap water and blood. *Environ Sci Technol* 36(8):1692-1698.
- Narotsky MG, Pegram RA, Kavlock RJ. 1997. Effect of dosing vehicle on the developmental toxicity of bromodichloromethane and carbon tetrachloride in rats. *Fundam Appl Toxicol* 40(1):30-36.
- Nieuwenhuijsen MJ, Grellier J, Iszatt N, Martinez D, Rahman MB, Villanueva CM. 2010. Literature Review of Meta-Analyses and Pooled Analyses of Disinfection By-Products in Drinking Water and Cancer and Reproductive Health Outcomes. In: *Contaminants of Emerging Concern in the Environment: Ecological and Human Health Considerations* (Halden RU, Ed). Washington DC: Amer Chemical Soc, 483-496.
- Nieuwenhuijsen MJ, Grellier J, Smith R, Iszatt N, Bennett J, Best N, et al. 2009. The epidemiology and possible mechanisms of disinfection by-products in drinking water. *Philos Transact a Math Phys Eng Sci* 367(1904):4043-4076.
- Nieuwenhuijsen MJ, Toledano MB, Elliott P. 2000. Uptake of chlorination disinfection by-products; a review and a discussion of its implications for exposure assessment in epidemiological studies. *J Expo Anal Environ Epidemiol* 10(6 Pt 1):586-599.
- Nuckols JR, Ashley DL, Lyu C, Gordon SM, Hinckley AF, Singer P. 2005. Influence of tap water quality and household water use activities on indoor air and internal dose levels of trihalomethanes. *Environ Health Perspect* 113(7):863-870.
- Patelarou E, Kargaki S, Stephanou EG, Nieuwenhuijsen M, Sourtzi P, Gracia E, et al. 2011. Exposure to brominated trihalomethanes in drinking water and reproductive outcomes. *Occup Environ Med* 68(6):438-445.
- Porter CK, Putnam SD, Hunting KL, Riddle MR. 2005. The effect of trihalomethane and haloacetic acid exposure on fetal growth in a Maryland county. *Am J Epidemiol* 162(4):334-344.
- Richardson SD, Plewa MJ, Wagner ED, Schoeny R, Demarini DM. 2007. Occurrence, genotoxicity, and carcinogenicity of regulated and emerging disinfection by-products in drinking water: a review and roadmap for research. *Mutat Res* 636(1-3):178-242.

- Riederer AM, Dhingra R, Blount BC, Steenland K. 2014. Predictors of blood trihalomethane concentrations in NHANES 1999-2006. *Environ Health Perspect* 122(7):695-702.
- Rivera-Nunez Z, Wright JM, Blount BC, Silva LK, Jones E, Chan RL, et al. 2012. Comparison of trihalomethanes in tap water and blood: a case study in the United States. *Environ Health Perspect* 120(5):661-667.
- Ruddick JA, Villeneuve DC, Chu I, Valli VE. 1983. A teratological assessment of four trihalomethanes in the rat. *J Environ Sci Health B* 18(3):333-349.
- Savitz DA, Singer PC, Herring AH, Hartmann KE, Weinberg HS, Makarushka C. 2006. Exposure to drinking water disinfection by-products and pregnancy loss. *Am J Epidemiol* 164(11):1043-1051.
- Shah PS. 2010. Parity and low birth weight and preterm birth: a systematic review and meta-analyses. *Acta Obstet Gynecol Scand* 89(7): 862-875.
- Silva LK, Backer LC, Ashley DL, Gordon SM, Brinkman MC, Nuckols JR, et al. 2013. The influence of physicochemical properties on the internal dose of trihalomethanes in humans following a controlled showering exposure. *J Expo Sci Environ Epidemiol* 23(1):39-45.
- Toledano MB, Nieuwenhuijsen MJ, Best N, Whitaker H, Hambly P, de Hoogh C, et al. 2005. Relation of trihalomethane concentrations in public water supplies to stillbirth and birth weight in three water regions in England. *Environ Health Perspect* 113(2):225-232.
- Weisel CP, Kim H, Haltmeier P, Klotz JB. 1999. Exposure estimates to disinfection by-products of chlorinated drinking water. *Environ Health Perspect* 107(2):103-110.
- Wright JM, Schwartz J, Dockery DW. 2003. Effect of trihalomethane exposure on fetal development. *Occup Environ Med* 60(3):173-180.
- Wright JM, Schwartz J, Dockery DW. 2004. The effect of disinfection by-products and mutagenic activity on birth weight and gestational duration. *Environ Health Perspect* 112(8):920-925.
- Zeng Q, Li M, Xie SH, Gu LJ, Yue J, Cao WC, et al. 2013. Baseline blood trihalomethanes, semen parameters and serum total testosterone: a cross-sectional study in China. *Environ Int* 54:134-140.
- Zeng Q, Chen YZ, Xu L, Chen HX, Luo Y, Li M, et al. 2014a. Evaluation of exposure to trihalomethanes in tap water and semen quality: a prospective study in Wuhan, China. *Reprod Toxicol* 46:56-63.
- Zeng Q, Zhou B, Cao WC, Wang YX, You L, Huang YH, et al. 2014b. Predictors of urinary trichloroacetic acid and baseline blood trihalomethanes concentrations among men in China. *Sci Total Environ* 493:806-811.
- Zhou WS, Xu L, Xie SH, Li YL, Li L, Zeng Q, et al. 2012. Decreased birth weight in relation to maternal urinary trichloroacetic acid levels. *Science of the Total Environment* 416:105-110.

**Table 1.** Maternal and neonatal infant characteristics in the study populations and according to birth outcome <sup>a</sup>

Variables	Study population [n (%)]	SGA [n (%)]	Mean birth weight (g)	Mean birth length (cm)	Mean gestational age (week)
<b>Total births</b>	1184 (100)	60 (5.1)	3321.8±404.1	50.0±1.0	38.7±4.6
<b>Season</b>					
Spring	393 (33.2)	14 (23.3)	3351.1±400.0	50.1±1.0	39.0±1.1
Summer	306 (25.8)	16 (26.7)	3313.0±407.7	50.0±1.3	39.1±1.3
Autumn	386 (32.6)	24 (40.0)	3312.3±409.1	50.1±0.8	39.1±1.4
Winter	99 (8.4)	6 (10.0)	3269.4±386.5	50.0±1.0	38.9±1.2
<b>Study city</b>					
Xiaogan	426 (36.0)	32 (53.3) <sup>**</sup>	3291.2±404.6 <sup>*</sup>	49.9±1.3 <sup>**</sup>	39.2±1.2 <sup>**</sup>
Wuhan	758 (64.0)	28 (46.7)	3338.9±403.0	50.1±0.9	39.0±1.3
<b>Infant's gender</b>					
Male	630 (53.2)	27 (45.0)	3357.0±403.6 <sup>**</sup>	50.2±0.9 <sup>**</sup>	39.0±1.1 <sup>*</sup>
Female	554 (46.8)	33 (55.0)	3282.9±401.3	49.9±1.2	39.1±1.4
<b>Weight gain during pregnancy</b>					
<15 kg	448 (38.9)	32 (55.2) <sup>*</sup>	3252.5±396.7 <sup>**</sup>	49.9±1.1 <sup>**</sup>	39.0±1.3
≥15 kg	705 (61.1)	26 (44.8)	3367.6±404.1	50.1±1.0	39.1±1.2
<b>Prenatal BMI</b>					
<28 kg/m <sup>2</sup>	811 (68.5)	46 (76.7)	3272.4±369.1 <sup>**</sup>	50.0±1.0 <sup>**</sup>	39.1±1.3
≥28 kg/m <sup>2</sup>	373 (31.5)	14 (23.3)	3429.1±453.5	50.2±1.2	39.0±1.2
<b>Education</b>					
Less than primary school	47 (4.0)	2 (3.3) <sup>*</sup>	3273.3±381.0	49.5±1.8 <sup>**</sup>	39.0±1.1
Junior and senior high school	624 (52.7)	41 (68.4)	3300.3±399.3	50.0±1.0	39.1±1.3
College and above	513 (43.3)	17 (28.3)	3352.3±410.5	50.2±0.9	39.0±1.2
<b>Household income, RMB, per month</b>					
<3000 Yuan	594 (50.2)	35 (58.3) <sup>*</sup>	3304.5±407.7	49.9±1.2	39.2±1.3
3000-<5000 Yuan	370 (31.3)	17 (28.3)	3316.7±382.9	50.1±0.8	39.0±1.2
≥5000 Yuan	220 (18.5)	8 (13.4)	3376.9±404.1	50.2±0.8	39.0±1.2
<b>Parity</b>					
No child	881 (74.4)	43 (71.7)	3316.6 ± 391.8	50.1 ± 1.0	39.1 ± 1.2
≥one child	303 (25.6)	17 (28.3)	3336.9 ± 438.1	50.0 ± 1.3	39.0 ± 1.4

<b>Usage of boiled water</b>					
Yes	1080 (91.2)	51 (85.0)	3324.7±403.9	50.0±1.1	39.0±1.3
No	104 (8.8)	9 (15.0)	3289.8±408.2	50.1±0.7	39.2±1.2
<b>Usage of filtered water</b>					
Yes	205 (17.4)	9 (15.0)	3294.4±401.1	50.1±1.0	39.1±1.1
No	975 (82.6)	51 (85.0)	3327.4±405.3	50.0±1.1	39.1±1.3
<b>Tap water consumption</b>					
<1200 mL/day	632 (55.8)	33 (55.0)	3294.4 ± 388.2	50.0 ± 1.1	39.2 ± 1.3 <sup>**</sup>
≥1200 mL/day	499 (44.2)	27 (45.0)	3349.7 ± 418.3	50.1 ± 1.0	39.0 ± 1.3
<b>Swim</b>					
Yes	7 (0.6)	0 (0.0)	3057.7 ± 347.6	49.8 ± 1.0	38.3 ± 1.3
No	1177 (99.4)	60 (100)	3323.1 ± 404.0	50.0 ± 1.0	39.1 ± 1.3
<b>Time of showering/bathing</b>					
<70 min/week	544 (48.1)	25 (46.3)	3322.4±394.1	50.1±0.9	39.1±1.3
≥70 min/week	586 (51.9)	29 (53.7)	3323.3±413.7	50.0±1.1	39.0±1.3
<b>Time of dishwashing</b>					
0 min/week	795 (73.0)	48 (81.3)	3325.4±412.3	50.1±1.1	39.0±1.6
<35 min/week	115 (10.6)	4 (6.8)	3276.4±383.8	50.1±0.7	39.1±1.4
≥35 min/week	179 (16.4)	7 (11.9)	3323.6±383.6	49.9±1.1	39.0±1.3
<b>Time of washing clothes</b>					
0 min/week	584 (53.4)	32 (56.2)	3333.7±426.4	50.1±1.1	39.0±1.2
<40 min/week	250 (22.9)	10 (17.5)	3322.9±386.4	50.1±0.9	39.0±1.3
≥40 min/week	259 (23.7)	15 (26.3)	3301.5±369.0	49.9±1.0	39.1±1.4
<b>Maternal age (years) (Mean)</b>	28.7±4.6	28.3±5.4	28.7±4.6	28.7±4.6	28.7±4.6 <sup>**</sup>

<sup>a</sup> 31 missing weight gain during pregnancy, 4 missing usage of filtered water, 53 missing tap water consumption, 54 missing time of showing/bathing, 95 missing time of washing dishes, 91 missing time of washing clothes.

\*  $p$ -value < 0.05, \*\*  $p$ -value < 0.01 for overall difference of fetal outcomes in the categorical and continuous variables.

**Table 2.** Distribution of maternal blood THM concentrations (ng/L) (n=1184)

Exposure variables <sup>a</sup>	Percent detected (95%CI)	Geometric mean (95%CI)	Median (95%CI)
TCM	92.5 (91.0, 94.0)	40.7 (38.0, 43.6)	50.7 (48.0, 53.0)
BDCM	57.4 (54.6, 60.2)	1.5 (1.4, 1.6)	2.5 (2.0, 2.9)
DBCM	33.5 (30.8, 36.2)	0.9 (0.9, 1.0)	0.5 (0.5, 0.5)
TBM	22.6 (20.2, 24.9)	1.6 (1.6, 1.6)	1.4 (1.4, 1.4)
Br-THMs <sup>b</sup>	-	5.3 (5.1, 5.5)	5.6 (5.2, 5.9)
TTHMs <sup>c</sup>	-	52.3 (49.8, 55.0)	57.7 (55.1, 59.9)

<sup>a</sup> The LODs for TCM, BDCM, DBCM and TBM were 1.9, 0.5, 0.7 and 2.0 ng/L, respectively. When the concentration was below the LOD, it was replaced with LOD/ $\sqrt{2}$ .

<sup>b</sup> Br-THMs: sum of BDCM, DBCM and TBM.

<sup>c</sup> TTHMs: sum of TCM and Br-THMs.



**Table 3.** Regression coefficients [ $\beta$  (95% CI)] for fetal development associated with categories of maternal blood THM concentrations (n=1184)

Blood THMs categories (ng/L)	Birth weight (g) <sup>a</sup>	Birth length (cm) <sup>a</sup>	Gestational age (week) <sup>b</sup>
<b>TCM</b>			
< 38.2	0	0	0
38.2 - 67.1	-25.90(-81.9, 30.13)	-0.08(-0.23, 0.07)	0.13(-0.05, 0.30)
>67.1	-48.23(-103.64, 7.19)	0.04(-0.11, 0.18)	0.15(-0.03, 0.32)
<i>p</i> for trend	0.08	0.63	0.10
<b>BDCM</b>			
<0.5	0	0	0
0.5-4.8	-27.41(-82.94, 28.11)	-0.03(-0.18, 0.12)	-0.05(-0.23, 0.12)
>4.8	-36.32(-91.22, 18.58)	-0.15(-0.29, -0.01)*	0.01(-0.16, 0.19)
<i>p</i> for trend	0.18	0.04	0.93
<b>DBCM</b>			
<0.7	0	0	0
0.7-2.6	4.66(-59.46, 68.78)	-0.05(-0.22, 0.12)	0.00(-0.20, 0.20)
>2.6	-4.98(-66.97, 57.02)	-0.20(-0.37, -0.04)**	0.04(-0.15, 0.23)
<i>p</i> for trend	0.92	0.02	0.72
<b>TBM</b>			
<2.0	0	0	0
2.0-2.4	19.01(-54.49, 92.51)	0.05(-0.15, 0.24)	-0.11(-0.34, 0.12)
>2.4	-24.72(-96.99, 47.55)	-0.06(-0.25, 0.14)	-0.06(-0.29, 0.17)
<i>p</i> for trend	0.66	0.72	0.44
<b>Br-THMs</b>			
<3.3	0	0	0
3.3-7.5	-12.99(-69.35, 43.36)	0.00(-0.15, 0.14)	-0.01(-0.19, 0.17)
>7.5	-25.53(-81.21, 30.15)	-0.04(-0.18, 0.11)	-0.08(-0.25, 0.10)
<i>p</i> for trend	0.37	0.62	0.39
<b>TTHMs</b>			
< 44.2	0	0	0
44.2-74.4	-59.09(-114.46, -3.71)*	-0.10(-0.25, 0.05)	0.06(-0.11, 0.23)
>74.4	-60.88(-116.18, -5.58)*	0.00(-0.15, 0.14)	0.14(-0.04, 0.31)
<i>p</i> for trend	0.03	0.96	0.12

<sup>a</sup> Adjusted for gestational age, infant's gender, maternal age, prenatal BMI, weight gain during pregnancy, education, parity and study city.

<sup>b</sup> Adjusted for infant's gender, maternal age, prenatal BMI, weight gain during pregnancy, household income, parity and study city.

\*  $p < 0.05$ , \*\*  $p < 0.01$

**Table 4.** ORs and 95% CIs for SGA with categories of maternal blood THM concentrations  
(n=1005)<sup>a, b</sup>

Blood THMs categories (ng/L)	SGA (n)	Adjusted OR (95% CI)
<b>TCM</b>		
<38.2	13	1
38.2 - 67.1	24	1.62(0.78, 3.36)
>67.1	23	1.48(0.71, 3.08)
<i>p</i> for trend		0.32
<b>BDCM</b>		
<0.5	25	1
0.5-4.8	18	1.30 (0.65, 2.59)
>4.8	17	1.15(0.58, 2.28)
<i>p</i> for trend		0.66
<b>DBCM</b>		
<0.7	40	1
0.7-2.6	10	1.16(0.54, 2.52)
>2.6	10	1.18(0.56, 2.49)
<i>p</i> for trend		0.62
<b>TBM<sup>c</sup></b>		
<2.0	52	1
≥2.0	8	0.81 (0.33, 1.99)
<i>p</i>		0.65
<b>Br-THMs</b>		
<3.3	16	1
3.3-7.5	18	1.48(0.71, 3.04)
>7.5	26	1.92(0.98, 3.79)
<i>p</i> for trend		0.06
<b>TTHMs</b>		
<44.2	9	1
44.2-74.4	28	2.91(1.32, 6.42) <sup>**</sup>
>74.4	23	2.25(1.01, 5.03) <sup>*</sup>
<i>p</i> for trend		0.08

<sup>a</sup> Excluding 179 large for gestational age (LGA) based on weight, defined as a live-born infant above the 10th percentile of birth weight for gestational age in a Chinese national referent population.

<sup>b</sup> Adjusted for infant's gender, maternal age, prenatal BMI, weight gain during pregnancy, education, household income, parity and study city.

<sup>c</sup> TBM was divided into two groups by LOD due to the small number of SGA in high level.

<sup>\*</sup>  $p < 0.05$ , <sup>\*\*</sup>  $p < 0.01$